

# 化学療法実施計画書（投与日程）

レジメン名=《1クール目》乳癌CAP+HER

(対象疾病= )

《初回開始日》 年 月 日

今回〔 〕クール目

身長= \_\_\_\_\_ cm

体重= \_\_\_\_\_ kg

体表面積= \_\_\_\_\_ m<sup>2</sup>

Ccr = \_\_\_\_\_

担当医	

薬剤 (商品名, 一般名)	1日投与量 (/m <sup>2</sup> or /kg)	投与法 点静・静注	投与時間 (min)	1クール投与スケジュール(日目)																																				
				1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30							
1 ハーセプチン(HER)《初回》	4mg/kg	点静	90	↓																																				
2 ハーセプチン《2回目以降》	2mg/kg	点静	60														↓																							
ゼローダ(CAP):A法 体表面積	1,650mg/m <sup>2</sup> *	経口		↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓		
★ゼローダ 1.31m <sup>2</sup> 未満	900mg × 2回/日																																							
1.31~1.64m <sup>2</sup> 未満	1,200mg × 2回/日																																							
1.64m <sup>2</sup> 以上	1,500mg × 2回/日																																							
前投薬																																								
投与日程																																								
副作用																																								
1クール期間	28日	クール回数																																						
休薬期間																																								
☆予想される重篤副作用	ハーセプチン ゼローダ	心障害, 24時間以内に現れるInfusion reaction 手足症候群(hand-foot syndrome)、肝障害																																						
☆副作用対策	ハーセプチン ゼローダ	心機能評価(心エコー、MUGAスキャン) ・初回は入院してInfusion reaction に関して観察する。 手足症候群対策: 保湿クリーム(1日5回)、ビタミンB6(60mg/day連日投与)、減量、休薬																																						
☆実施中止条件	生化学項目 副作用 その他																																							